

On the Isolation of the Roots of a Non-Linear Operator*

A. A. GOLDSTEIN

Department of Mathematics, University of Washington, Seattle, Washington 98105

F. C. JOHNSON

Boeing Scientific Research Laboratories, Seattle

AND

JURGEN SPIESS

University of Hamburg, Hamburg

Received July 9, 1969

Certain problems in optimal control may be formulated by the theory of Lagrange multipliers. Stationary controls are thus roots of an operator equation. See, e.g., Ref. [1, 2]. We give below a measure of the isolation of these roots.

We will follow the notational convention that $f'(x)$ represents the Gateaux derivative of f evaluated at x , and that $f'(x, h)$ is the differential of f at x in the direction h .

THEOREM. *Let G be a Gateaux differentiable mapping from an open convex subset C of a normed linear space N to a Hilbert space E . Assume that G has a root z in C , that G' satisfies $\|G'(z, x - z)\| \geq \mu \|x - z\|$ and $\|G'(z, x - z) - G'(y, x - z)\| \leq L \|x - z\| \|y - z\|$ for all $x, y \in S$ and some positive μ and L . Here S denotes the set $\{x \in C: \|x - z\| < \mu/L\}$. Then z is the only root of G in S .*

As an aside, we observe that since G is actually Frechet differentiable at z , the root z can easily be seen to be isolated.

Proof. Let $\phi(x) = \frac{1}{2} \langle G(x), G(x) \rangle$. We have $\phi(z) = 0 = \phi'(z)$ and $\phi(x) = \int_0^1 \phi'(z + t(x - z), x - z) dt$.

We show that $\phi(x) > 0$ for all $x \in S - \{z\}$.

* This research was supported in part by the Air Force Office of Scientific Research and by the Boeing Scientific Research Laboratories.

Let $h = x - z$ and $y = z + th$ with $0 < t \leq 1$; then

$$\begin{aligned}
 \phi'(y, h) &= \langle G(y), G'(y, h) \rangle \\
 &= \langle G'(y, h), G'(y, h) \rangle + \langle G(y) - G(z) - G'(z, h), G'(y, h) \rangle \\
 &\quad + \langle G'(z, h) - G'(y, h), G'(y, h) \rangle \\
 &\geq \{ \|G'(y, h)\| - \|G(y) - G(z) - G'(z, h)\| \\
 &\quad - \|G'(z, h) - G'(y, h)\| \} \|G'(y, h)\|.
 \end{aligned} \tag{3}$$

We now estimate:

(a) Choose $f \in E^*$, such that $\|f\| = 1$ and

$$\begin{aligned}
 &\|G(y) - G(z) - G'(z, h)\| \\
 &= f(G(y) - G(z) - G'(z, h)) = f(G(y)) - f(G(z)) - f(G'(z, h)) \\
 &= f(G'(\xi, h)) - f(G'(z, h)) = f(G'(\xi, h) - G'(z, h)) \\
 &\leq \|G'(\xi, h) - G'(z, h)\| \leq L \|\xi - z\| \|h\| \leq Lt \|h\|^2.
 \end{aligned}$$

In the above, $\xi = z + \tau h$ and $0 < \tau < t$.

(b) $\|G'(z, h) - G'(y, h)\| \leq L \|z - y\| \|h\| = Lt \|h\|^2$.

(c) Since $\|G'(z, h)\| - \|G'(y, h)\| \leq \|G'(z, h) - G'(y, h)\| \leq Lt \|h\|^2$, we have that

$$\|G'(y, h)\| \geq \|G'(z, h)\| - Lt \|h\|^2 \geq (\mu - tL \|h\|) \|h\|.$$

Combining a, b, c, in (3), we get

$$\phi'(y, h) \geq \|h\|^2 (\mu - t\alpha)(\mu - 3t\alpha) \quad (\alpha = L \cdot \|h\|);$$

hence

$$\begin{aligned}
 \phi(x) &= \int_0^1 \phi'(y, h) dt \geq \|h\|^2 \int_0^1 (\mu - t\alpha)(\mu - 3t\alpha) dt = \|h\|^2 \cdot (\mu - \alpha)^2 \\
 &= \|x - z\|^2 \cdot (\mu - L \|x - z\|)^2.
 \end{aligned}$$

Thus $\phi(x) > 0$, if $x \in S - \{z\}$.

LEMMA. Let H_1 and H_2 be Hilbert spaces and A be a bounded linear operator from H_1 onto H_2 . Then AA^* has a bounded inverse.

Proof. We first observe that the null space of A , $N(A)$, equals $\{0\}$. Take $l \in H_2$, $l \neq 0$, and set $y = A^*l$. Suppose that $y \in N(A)$ and $y \neq 0$. Since $l \neq 0$, there exists $x \in N^\perp(A)$ such that $Ax = l$. Thus we have

$$0 = \langle x, y \rangle = \langle x, A^*l \rangle = \langle Ax, l \rangle = \langle l, l \rangle,$$

a contradiction. Moreover, we have $N(A^*) = \{0\}$ and therefore $N(AA^*) = \{0\}$. Thus, AA^* is a one-to-one map. Since the range of A is closed, the range of A^* is closed [3, p. 488]. Therefore, the range of AA^* is closed. Suppose that the range R of AA^* is not H_2 , and take $l \in R^\perp$, $l \neq 0$. Then $0 = \langle l, AA^*l \rangle = \langle A^*l, A^*l \rangle$, a contradiction. Therefore AA^* is onto, and by the open mapping theorem $(AA^*)^{-1}$ is bounded.

Remark. If $B: H_1 \rightarrow H_1$ and B has a bounded inverse, then ABA^* has a bounded inverse.

Application to Constrained Minimization

Let A be an operator from H_1 to H_2 and f a functional defined on H_1 . We consider the problem of minimizing f on the set $S = \{x \in H_1 : A(x) = 0\}$.

Let z be a stationary point of f on S , i.e., $z \in S$, the Fréchet derivatives A' and f' exist at z and there exists $\Lambda \in H_2$ such that $f'(z, h) = \langle \Lambda, A'(z, h) \rangle$ for all $h \in H_1$. Let C_0 be a convex neighborhood of z and assume that there exist positive numbers $\alpha, \beta, \gamma, \psi, \Omega, \theta$ and ν such that the following conditions hold on C_0 :

(i) f has a second Gateaux derivative which satisfies

$$\|f''(x) - f''(z)\| \leq \alpha \|x - z\|.$$

(ii) A has a second Gateaux derivative which satisfies

$$\|A''(x) - A''(z)\| \leq \beta \|x - z\|.$$

(iii) $\|A''(x)\| \leq \gamma$

(iv) Set $D(x) = A'(x) A'^*(x)$. Then $D^{-1}(x)$ exists and satisfies

$$\|D^{-1}(x) - D^{-1}(z)\| \leq \psi \|x - z\|$$

(v) $\|D^{-1}(x)\| \leq \Omega$.

(vi) $\|A'(x)\| \leq \theta$.

(vii) $\|f'(x) - f'(z)\| \leq \nu \|x - z\|$.

Let $H = H_1^* \oplus H_2$ and $S_0 = C_0 \oplus H_2$. Let G be a mapping from S_0 to H defined by

$$G(p) = (f'(x) - \lambda \circ A'(x), A(x)),$$

where $p = (x, \lambda)$ and $\lambda \circ A'(x) \in H_1^*$ is defined by $(\lambda \circ A'(x))h = \langle A'(x, h), \lambda \rangle$. Note that a zero of G is a stationary point of f on S and conversely. Let $p_0 = (z, \Lambda)$ and define the operator $C \in B(H_1, H_1)$ by

$$Ch = f''(z, h) - \Lambda \circ A''(z, h).$$

We shall show that $G'(p_0)^{-1}$ is bounded. Let $\eta = \|G'(p_0)^{-1}\|$ and define

$$\begin{aligned} L &= \alpha + \beta \|A\| + 3\gamma, \\ M &= \nu\Omega\Theta + (\psi \|A'(z)\| + \gamma\Omega) \|f(z)\|. \end{aligned}$$

THEOREM. *Assume that $A'(z)$ maps H_1 onto H_2 and that C^{-1} exists. Let N be a sphere centered at z with radius $r = 1/(1+M)\eta L$. Then z is a unique stationary point of f with respect to S on the set $N \cap C_0$.*

Proof. Let $\Delta p = (h, l)$. Then

$$G'(p, \Delta p) = (f''(x, h) - \lambda \circ A''(x, h) - l \circ A'(x), A'(x, h)).$$

We note first that $G'(p_0)$ maps H onto H . For, let $(a, b) \in H$. Then we seek $\Delta p \in H$ such that $G'(p_0, \Delta p) = (a, b)$. Since the operator $\lambda \circ A'(z)$ also has the representation $A'^*(z)\lambda$, then, dropping the z argument for simplicity, we are led to the equations

$$\begin{aligned} Ch - A'^*l &= a, \\ A'h &= b, \end{aligned}$$

which has the solution

$$\begin{aligned} l &= (A'C^{-1}A'^*)^{-1}(b - A'C^{-1}a), \\ h &= C^{-1}(a + A'^*l), \end{aligned}$$

where the existence of $(A'C^{-1}A'^*)^{-1}$ is established by the remark following the lemma.

It follows that $G'(p_0)$ is also one-to-one and the open mapping theorem implies that $G'(p_0)^{-1}$ is bounded. Since we have defined $\eta = \|G'(p_0)^{-1}\|$, we have that

$$\|G'(p_0, \Delta p)\| \geq \frac{1}{\eta} \|\Delta p\|.$$

A straightforward computation shows that G' is Lipschitz continuous at p_0 . We have

$$\begin{aligned} \|G'(p_0, \Delta p) - G'(p, \Delta p)\| &\leq (\alpha + \beta \|A\| + 2\gamma) \|x - z\| \|\Delta p\| + \gamma \|A - \lambda\| \|\Delta p\| \\ &\quad - L \|p - p_0\| \|\Delta p\|. \end{aligned}$$

Therefore, G' is Lipschitz continuous at p_0 with constant L . It follows then from the theorem that if $\|p - p_0\| \leq 1/\eta L$ and $p \neq p_0$, then $G(p) \neq 0$.

Let $y \in C_0$ such that y is a stationary point of f on S , and let κ be the corresponding multiplier. Then (y, κ) satisfies

$$f'(y) = A'^*(y)\kappa,$$

and it follows from Lemma 1 that

$$\kappa = D^{-1}(y) A'(y) f'(y).$$

We have then

$$\begin{aligned} \|A - \kappa\| &= \|D^{-1}(z) A'(z) f'(z) - D^{-1}(y) A'(y) f'(y)\| \\ &\leq (\psi \|A'(z)\| \|f(z)\| + \gamma \Omega \|f(z)\| + \nu \Omega \theta) \|y - z\| \\ &= M \|y - z\|. \end{aligned}$$

Thus, we have that

$$\|p_0 - (y, \kappa)\| \leq (1 + M) \|y - z\|.$$

Hence, if $y \in N \cap C_0$, then $\|p_0 - (y, \kappa)\| \leq 1/L\eta$, and since G has a unique root on $N \cap C_0$, we must have $(y, K) = p_0$. Q.E.D.

REFERENCES

1. A. A. GOLDSTEIN AND J. S. MEDITCH, Non-linear optimal controls, *J. Comput. System Sci.* **4** (1970), 29-37.
2. F. C. JOHNSON, Norm minimization on nonlinear manifolds in Hilbert space, Boeing document D1-82-0758, 1968.
3. N. DUNFORD AND J. T. SCHWARTZ, "Linear Operators I," Interscience, New York, 1966.